

THE COHOMOLOGY RING OF THE GKM GRAPH OF A FLAG MANIFOLD

大阪市立大学 大学院理学研究科 福川由貴子 (Yukiko Fukukawa)
Department of Mathematics,
Osaka City University

1. INTRODUCTION

Let T be a torus of dimension n and M a closed smooth T -manifold. The equivariant cohomology of M , denoted $H_T^*(M)$, contains a lot of geometrical information on M . Moreover it is often easier to compute $H_T^*(M)$ than $H^*(M)$ by virtue of the Localization Theorem which implies that the restriction map

$$(1.1) \quad \iota^*: H_T^*(M) \rightarrow H_T^*(M^T)$$

to the T -fixed point set M^T is often injective, in fact, this is the case when $H^{odd}(M) = 0$. When M^T is isolated, $H_T^*(M^T) = \bigoplus_{p \in M^T} H_T^*(p)$ and hence $H_T^*(M^T)$ is a direct sum of copies of a polynomial ring in n variables because $H_T^*(p) = H^*(BT)$.

Therefore we are in a nice situation when $H^{odd}(M) = 0$ and M^T is isolated. Goresky-Kottwitz-MacPherson [2] (see also [3, Chapter 11]) found that under the further condition that the weights at a tangential T -module are pairwise linearly independent at each $p \in M^T$, the image of ι^* in (1.1) above is determined by the fixed point sets of codimension one subtori of T when \mathbb{Q} is tensored in cohomology. Their result motivated Guillemin-Zara [4] to associate a labeled graph \mathcal{G}_M with M and define the ‘‘cohomology’’ ring $\mathcal{H}^*(\mathcal{G}_M)$ of \mathcal{G}_M , which is a subring of $\bigoplus_{p \in M^T} H^*(BT)$. Then the result of Goresky-Kottwitz-MacPherson can be stated that $H_T^*(M) \otimes \mathbb{Q}$ is isomorphic to $\mathcal{H}^*(\mathcal{G}_M) \otimes \mathbb{Q}$ as graded rings when M satisfies the conditions mentioned above.

The result of Goresky-Kottwitz-MacPherson can be applied to many important T -manifolds M such as flag manifolds and compact smooth toric varieties etc. When M is such a nice manifold, $H_T^*(M)$ is often known to be isomorphic to $\mathcal{H}^*(\mathcal{G}_M)$ without tensoring with \mathbb{Q} (see [1], [5], [6] for example). We determine the ring structure of $\mathcal{H}^*(\mathcal{G}_M)$ or $\mathcal{H}^*(\mathcal{G}_M) \otimes \mathbb{Z}[\frac{1}{2}]$ when M is a flag manifold of classical type directly without using the fact

that $H_T^*(M)$ is isomorphic to $\mathcal{H}^*(\mathcal{G}_M)$ ([7]). In my talk, I introduced the result when M is a flag manifold of type A. This is a joint work with Hiroaki Ishida and Mikiya Masuda and the details can be found in [7].

2. LABELED GRAPH AND ITS COHOMOLOGY FOR TYPE A_{n-1}

Let $\{t_i\}_{i=1}^n$ be a basis of $H^2(BT)$, so that $H^*(BT)$ can be identified with a polynomial ring $\mathbb{Z}[t_1, t_2, \dots, t_n]$. We take an inner product on $H^2(BT)$ such that the basis $\{t_i\}$ is orthonormal. Then

$$(2.1) \quad \Phi(A_{n-1}) := \{\pm(t_i - t_j) \mid 1 \leq i < j \leq n\}$$

is a root system of type A_{n-1} .

Definition. The labeled graph associated with $\Phi(A_{n-1})$, denoted \mathcal{A}_n , is a graph with labeling ℓ defined as follows.

- The vertex set of \mathcal{A}_n is the permutation group S_n on $\{1, 2, \dots, n\}$.
- Two vertices w, w' in \mathcal{A}_n are connected by an edge $e_{w, w'}$ if and only if there is a transposition $(i, j) \in S_n$ such that $w' = w(i, j)$, in other words,

$$w'(i) = w(j), \quad w'(j) = w(i) \quad \text{and} \quad w'(r) = w(r) \quad \text{for } r \neq i, j.$$
- The edge $e_{w, w'}$ is labeled by $\ell(e_{w, w'}) := t_{w(i)} - t_{w'(i)}$.

Definition. The cohomology ring of \mathcal{A}_n , denoted $\mathcal{H}^*(\mathcal{A}_n)$, is defined to be the subring of $\text{Map}(V(\mathcal{A}_n), H^*(BT)) = \bigoplus_{v \in V(\mathcal{A}_n)} H^*(BT)$, where $V(\mathcal{A}_n)$ denotes the set of vertices of \mathcal{A}_n , i.e. $V(\mathcal{A}_n) = S_n$, satisfying the following condition:


$f \in \text{Map}(V(\mathcal{A}_n), H^*(BT))$ is an element of $\mathcal{H}^*(\mathcal{A}_n)$ if and only if $f(v) - f(v')$ is divisible by $\ell(e)$ in $H^*(BT)$ whenever the vertices v and v' are connected by an edge e in \mathcal{A}_n .

For each $i = 1, \dots, n$, we define elements τ_i, t_i of $\text{Map}(V(\mathcal{A}_n), H^*(BT))$ by

$$(2.2) \quad \tau_i(w) := t_{w(i)}, \quad t_i(w) := t_i \quad \text{for } w \in S_n.$$

In fact, both τ_i and t_i are elements of $\mathcal{H}^2(\mathcal{A}_n)$.

Example. The case $n = 3$. The root system $\Phi(A_2)$ is $\{\pm(t_i - t_j) \mid 1 \leq i < j \leq 3\}$. The labeled graph \mathcal{A}_3 and τ_i for $i = 1, 2, 3$ are as follows.



Theorem 2.1. Let \mathcal{A}_n be the labeled graph associated with the root system $\Phi(A_{n-1})$ of type A_{n-1} in (2.1). Then

$$\mathcal{H}^*(\mathcal{A}_n) = \mathbb{Z}[\tau_1, \dots, \tau_n, t_1, \dots, t_n] / (e_i(\tau) - e_i(t) \mid i = 1, \dots, n),$$

where $e_i(\tau)$ (resp. $e_i(t)$) is the i^{th} elementary symmetric polynomial in τ_1, \dots, τ_n (resp. t_1, \dots, t_n).

To prove this theorem, we need the following two lemmas.

Lemma 2.2. $\mathcal{H}^*(\mathcal{A}_n)$ is generated by $\tau_1, \dots, \tau_n, t_1, \dots, t_n$ as a ring.

Proof. We shall prove the lemma by induction on n . When $n = 1$, $\mathcal{H}^*(\mathcal{A}_1)$ is generated by t_1 since \mathcal{A}_1 is a point; so the lemma holds.

Suppose that the lemma holds for $n - 1$. Then it suffices to show that any homogenous element f of $\mathcal{H}^*(\mathcal{A}_n)$, say of degree $2k$, can be expressed as a polynomial in τ_i 's and t_i 's. For each $i = 1, \dots, n$, we set

$$V_i := \{w \in S_n \mid w(i) = n\}$$

and consider the labeled full subgraph \mathcal{L}_i of \mathcal{A}_n with V_i as the vertex set. Note that \mathcal{L}_i can naturally be identified with \mathcal{A}_{n-1} for any i .

Let

$$(2.3) \quad 1 \leq q \leq \min\{k + 1, n\}$$

and assume that

$$(2.4) \quad f(v) = 0 \quad \text{for any } v \in V_i \text{ whenever } i < q.$$

A vertex w in V_q is connected by an edge in \mathcal{A}_n to a vertex v in V_i if and only if $v = w(i, q)$. In this case $f(w) - f(v)$ is divisible by $t_{w(i)} - t_{w(q)} = t_{w(i)} - t_n$ and $f(v) = 0$ whenever $i < q$ by (2.4), so $f(w)$ is divisible by $t_{w(i)} - t_n$ for $i < q$. Thus, for each $w \in V_q$, there is an element $g^q(w) \in \mathbb{Z}[t_1, \dots, t_n]$ such that

$$(2.5) \quad f(w) = (t_{w(1)} - t_n)(t_{w(2)} - t_n) \dots (t_{w(q-1)} - t_n)g^q(w)$$

where $g^q(w)$ is homogenous and of degree $2(k + 1 - q)$ because $f(w)$ is homogenous and of degree $2k$.

One expresses

$$(2.6) \quad g^q(w) = \sum_{r=0}^{k+1-q} g_r^q(w)t_n^r$$

with homogenous polynomials $g_r^q(w)$ of degree $2(k + 1 - q - r)$ in $\mathbb{Z}[t_1, \dots, t_{n-1}]$. Then there is a polynomial G_r^q in τ_i 's (except τ_q) and t_i 's (except t_n) such that $G_r^q(w) = g_r^q(w)$ for any $w \in V_q$, because g_r^q restricted to \mathcal{L}_q is an element of $\mathcal{H}^*(\mathcal{L}_q) = \mathcal{H}^*(\mathcal{A}_{n-1})$.

Since $\tau_i(w) = t_{w(i)}$ and $w(i) = n$ for $w \in V_i$, we have

$$(2.7) \quad \prod_{i=1}^{q-1} (\tau_i - t_n)(w) = 0 \quad \text{for any } w \in V_i \text{ whenever } i < q.$$

Therefore, it follows from (2.5), (2.6), the Claim above and (2.7) that putting $G^q = \sum_{r=0}^{k+1-q} G_r^q t_n^r$, we have

$$\begin{aligned} (f - G^q \prod_{i=1}^{q-1} (\tau_i - t_n))(w) &= f(w) - g^q(w) \prod_{i=1}^{q-1} (t_{w(i)} - t_n) \\ &= 0 \quad \text{for any } w \in V_i \text{ whenever } i \leq q. \end{aligned}$$

Therefore, subtracting the polynomial $G^q \prod_{i=1}^{q-1} (\tau_i - t_n)$ from f , we may assume that

$$f(v) = 0 \quad \text{for any } v \in V_i \text{ whenever } i < q + 1.$$

The above argument implies that f finally takes zero on all vertices of \mathcal{A}_n (which means $f = 0$) by subtracting a polynomial in τ_i 's and t_i 's, and this completes the induction step. \square

We abbreviate the polynomial ring $\mathbb{Z}[\tau_1, \dots, \tau_n, t_1, \dots, t_n]$ as $\mathbb{Z}[\tau, t]$. The canonical map $\mathbb{Z}[\tau, t] \rightarrow \mathcal{H}^*(\mathcal{A}_n)$ is a grade preserving homomorphism which is surjective by Lemma 2.2. Let $e_i(\tau)$ (resp. $e_i(t)$) denote the i^{th} elementary symmetric polynomial in τ_1, \dots, τ_n (resp. t_1, \dots, t_n). It easily follows from (2.2) that $e_i(\tau) = e_i(t)$ for $i = 1, \dots, n$. Therefore the canonical map above induces a grade preserving epimorphism

$$(2.8) \quad \mathbb{Z}[\tau, t]/(e_1(\tau) - e_1(t), \dots, e_n(\tau) - e_n(t)) \rightarrow \mathcal{H}^*(\mathcal{A}_n).$$

Remember that the Hilbert series of a graded ring $A^* = \bigoplus_{j=0}^{\infty} A^j$, where A^j is the degree j part of A^* and of finite rank over \mathbb{Z} , is a formal power series defined by

$$F(A^*, s) := \sum_{j=0}^{\infty} (\text{rank}_{\mathbb{Z}} A^j) s^j.$$

In order to prove that the epimorphism in (2.8) is an isomorphism, it suffices to verify the following lemma because the modules in (2.8) are both torsion free.

Lemma 2.3. *The Hilbert series of the both sides at (2.8) coincide, in fact, they are given by $\frac{1}{(1-s^2)^n} \prod_{i=1}^n (1 - s^{2i})$.*

Proof. (1) Calculation of LHS at (2.8). Let $e_i := e_i(\tau) - e_i(t)$. It follows from the exact sequence

$$0 \rightarrow (e_1, \dots, e_n) \rightarrow \mathbb{Z}[\tau, t] \rightarrow \mathbb{Z}[\tau, t]/(e_1, \dots, e_n) \rightarrow 0$$

that we have

$$(2.9) \quad F(\mathbb{Z}[\tau, t]/(e_1, \dots, e_n), s) = F(\mathbb{Z}[\tau, t], s) - F((e_1, \dots, e_n), s).$$

Here, since $\deg \tau_i = \deg t_i = 2$, we have

$$(2.10) \quad F(\mathbb{Z}[\tau, t], s) = \frac{1}{(1-s^2)^{2n}}$$

as easily checked; so it suffices to calculate $F((e_1, \dots, e_n), s)$.

For $I \subset [n]$ we set $e_I := \prod_{i \in I} e_i$. Then it follows from the Inclusion-Exclusion principle that

$$(2.11) \quad F((e_1, \dots, e_n), s) = \sum_{\emptyset \neq I \subset [n]} (-1)^{|I|-1} F((e_I), s)$$

and since $F((e_I), s) = s^{\deg e_I} / (1-s^2)^{2n}$ and $\deg e_I = \sum_{i \in I} 2i$, it follows from (2.11) that

$$(2.12) \quad F((e_1, \dots, e_n), s) = \sum_{\emptyset \neq I \subset [n]} (-1)^{|I|-1} \frac{s^{\sum_{i \in I} 2i}}{(1-s^2)^{2n}}.$$

Therefore it follows from (2.9), (2.10) and (2.12) that

$$(2.13) \quad \begin{aligned} F(\mathbb{Z}[\tau, t]/(e_1, \dots, e_n), s) &= \frac{1}{(1-s^2)^{2n}} - \sum_{\emptyset \neq I \subset [n]} (-1)^{|I|-1} \frac{s^{\sum_{i \in I} 2i}}{(1-s^2)^{2n}} \\ &= \frac{1}{(1-s^2)^{2n}} \sum_{I \subset [n]} (-1)^{|I|} s^{\sum_{i \in I} 2i} \\ &= \frac{1}{(1-s^2)^{2n}} \prod_{i=1}^n (1-s^{2i}). \end{aligned}$$

(2) Calculation of RHS at (2.8). Let $d_n(k) := \text{rank}_{\mathbb{Z}} \mathcal{H}^{2k}(\mathcal{A}_n)$. Then

$$(2.14) \quad F(\mathcal{H}^*(\mathcal{A}_n), s) = \sum_{k=0}^{\infty} d_n(k) s^{2k}.$$

Recall the argument in the proof of Lemma 2.2. Since g_r^q in (2.6) belongs to $\mathcal{H}^{2(k+1-q-r)}(\mathcal{L}_q) = \mathcal{H}^{2(k+1-q-r)}(\mathcal{A}_{n-1})$ as shown in the Claim there, the rank of the module consisting of those g^q in (2.5) and (2.6) is given by

$$\sum_{r=0}^{k+1-q} d_{n-1}(k+1-q-r) = \sum_{r=0}^{k+1-q} d_{n-1}(r).$$

Therefore, noting (2.3), we see that the argument in the proof of Lemma 2.2 implies

$$d_n(k) = \sum_{q=1}^{\min\{k+1, n\}} \sum_{r=0}^{k+1-q} d_{n-1}(r),$$

in other words, if we set $d_{n-1}(j) = 0$ for $j < 0$, then

(2.15)

$$d_n(k) = \begin{cases} \sum_{i=1}^n id_{n-1}(k+1-i) & \text{if } k \leq n-1, \\ \sum_{i=1}^n id_{n-1}(k+1-i) + n \sum_{i=n+1}^{k+1} d_{n-1}(k+1-i) & \text{if } k \geq n. \end{cases}$$

We shall abbreviate $F(\mathcal{H}^*(\mathcal{A}_n), s)$ as $F_n(s)$. Then, plugging (2.15) in (2.14), we obtain

$$\begin{aligned} F_n(s) &= \sum_{k=0}^{\infty} \left(d_{n-1}(k) + 2d_{n-1}(k-1) + \dots + nd_{n-1}(k+1-n) \right) s^{2k} \\ &\quad + n \sum_{k=n}^{\infty} \left(d_{n-1}(k-n) + \dots + d_{n-1}(1) + d_{n-1}(0) \right) s^{2k} \\ &= F_{n-1}(s) + 2s^2 F_{n-1}(s) + \dots + ns^{2n-2} F_{n-1}(s) \\ &\quad + n \left(d_{n-1}(0) s^{2n} \frac{1}{1-s^2} + d_{n-1}(1) s^{2n+2} \frac{1}{1-s^2} + \dots \right) \\ &= F_{n-1}(s) \left(1 + 2s^2 + \dots + ns^{2n-2} \right) + n \frac{s^{2n}}{1-s^2} F_{n-1}(s) \\ &= \frac{1-s^{2n}}{1-s^2} F_{n-1}(s). \end{aligned}$$

On the other hand, $F_1(s) = 1/(1-s^2)$ since $\mathcal{H}^*(\mathcal{A}_1) = \mathbb{Z}[t_1]$. It follows that

$$F_n(s) = \frac{1}{(1-s^2)^{2n}} \prod_{i=1}^n (1-s^{2i}).$$

This together with (2.13) proves the lemma. \square

REFERENCES

- [1] A. Arabia, *Cohomologie T-équivariante de G/B pour un groupe G de Kac-Moody*, C. R. Acad. Sci. Paris Sér. I. Math. 302 (1986) 631–634.
- [2] M. Goresky, R. Kottwitz and R. MacPherson, *Equivariant cohomology, Koszul duality and the localisation theorem*, Invent. Math. 131 (1998) 25–83.
- [3] V. Guillemin and S. Sternberg, *Supersymmetry and Equivariant de Rham Theory*, Springer, Berlin–Heidelberg–New York, 1999.
- [4] V. Guillemin and C. Zara, *1-skeleta, Betti numbers and equivariant cohomology*, Duke Math. J. 107 (2001) 283–349.
- [5] M. Harada, A. Henriques and T. Holm, *Computation of generalized equivariant cohomologies of Kac-Moody flag varieties*, Adv. in Math. 197 (2005), 198–221.
- [6] H. Maeda, M. Masuda and T. Panov, *Torus graphs and simplicial posets*, Adv. Math. 212 (2007), 458–483.
- [7] Y. Fukukawa, H. Ishida and M. Masuda, *The cohomology ring of the GKM graph of a flag manifold of classical type*, preprint.