

Squeezing on a Certain \mathbb{L} -space

岡山理科大学・理学部 山崎 正之 (Masayuki Yamasaki) ¹

Faculty of Science, Okayama University of Science

1. INTRODUCTION

In a joint 2006 paper [2], E. Pedersen and I proved a certain stability result for controlled L -groups. The proof depended on a construction called the Alexander trick. In this note I describe a modified Alexander trick which can be used to give a built-in squeezing mechanism of a certain \mathbb{L} -space. This should replace the “barycentric subdivision argument” used in [4].

2. ITERATED MAPPING CYLINDERS

Let X be a finite polyhedron, and M be a topological space. We are interested in a map $p : M \rightarrow X$ which has an iterated mapping cylinder decomposition in the sense of Hatcher [1]: there is a partial order on the set of the vertices of X such that, for each simplex Δ of X ,

- (1) the partial order restricts to a total order of the vertices of Δ

$$v_0 < v_1 < \cdots < v_n ,$$

- (2) $p^{-1}(\Delta)$ is the iterated mapping cylinder of a sequence of maps

$$F_{v_0} \longrightarrow F_{v_1} \longrightarrow \cdots \longrightarrow F_{v_n} ,$$

- (3) the restriction $p|_{p^{-1}(\Delta)}$ is the natural map induced from the iterated mapping cylinder structure of $p^{-1}(\Delta)$ above and the iterated mapping cylinder structure of Δ coming from the sequence

$$\{v_0\} \longrightarrow \{v_1\} \longrightarrow \cdots \longrightarrow \{v_n\} .$$

To simplify the situation we assume that X is an n -simplex Δ with vertices v_0, v_1, \dots, v_n . The edge $|v_0, v_1|$ is the mapping cylinder $v_0 \times \{0 \leq t_1 \leq 1\} / (v_0, 1) \sim v_1$, the face $|v_0, v_1, v_2|$ is the mapping cylinder $|v_0, v_1| \times \{0 \leq t_2 \leq 1\} / (x, 1) \sim v_2, \dots$, and $\Delta = |v_0, \dots, v_n|$ is the mapping cylinder $|v_0, \dots, v_{n-1}| \times \{0 \leq t_n \leq 1\} / (x, 1) \sim v_n$. Thus we can assign a point in Δ to each $(t_1, \dots, t_n) \in [0, 1]^n$. (t_1, \dots, t_n) is *pseudo-coordinates* of the point in the sense that the coordinates are not uniquely determined

¹This work was supported by KAKENHI 20540100.

by the point. If $(\lambda_0, \dots, \lambda_n)$ are the barycentric coordinates of a point $x \in \Delta$, *i.e.* $x = \sum \lambda_i v_i$ ($\lambda_0 + \dots + \lambda_n = 1$), then t_i is equal to $\lambda_i / (\lambda_0 + \dots + \lambda_i)$, when defined, and is indeterminate when $\lambda_0 = \dots = \lambda_i = 0$.

For each vertex v of Δ , define a simplicial map $s^v : \Delta \rightarrow \Delta$ by:

$$s^v(u) = \begin{cases} v & \text{for a vertex } u \text{ with } u < v, \\ u & \text{for a vertex } u \text{ with } u \geq v. \end{cases}$$

For example, s^{v_0} is the identity map, and s^{v_n} is the constant map which sends every point of Δ to v_n . A strong deformation retraction $s_t^v : \Delta \rightarrow \Delta$ is defined by $s_t^v(x) = (1-t)x + t s^v(x)$, where $x \in \Delta$ and $t \in [0, 1]$. Note that this strong deformation retraction s_t^v is covered by a deformation \tilde{s}_t^v on M , since M has an iterated mapping cylinder structure. Also note that $s_t^{v_j}$ ($t > 0$) changes the t_j pseudo-coordinate but fixes the other pseudo-coordinates t_i ($i \neq j$).

3. ALEXANDER TRICKS

Let M be an iterated mapping cylinder of maps

$$F_{v_0} \longrightarrow F_{v_1} \longrightarrow \dots \longrightarrow F_{v_n},$$

and $p : M \rightarrow \Delta = |v_0, \dots, v_n|$ be the projection from M to the ordered n -simplex Δ as in the previous section. Suppose c is a quadratic Poincaré $(n+2)$ -ad on $p : M \rightarrow \Delta$, such that $\partial_i c$ is a quadratic Poincaré $(n+1)$ -ad on $p|_{p^{-1}(\partial_i \Delta)}$, $i = 0, \dots, n$ ([4] [5]). Such an $(n+2)$ -ad c is said to be *proper on Δ* or simply *proper*.

We will describe a version of Alexander trick for such a proper $(n+2)$ -ad c . First fix a positive integer N (“*height*”) and pick up a vertex $v = v_j$ of Δ toward which we try to squeeze the objects. Triangulate the closed interval $I_N = [0, N]$ using unit intervals and represent each simplex by its barycenter. Use these points to construct the symmetric Poincaré triad e of $(I_N; 0, N)$. Take the tensor product of c and e and denote it by $c \times I_N$. This is a geometric object on $M \times I_N$ which gives a cobordism between $c \times 0$ and the $(n+2)$ -ad c' defined by:

$$c' = c \times N \cup \partial_j c \times I_N,$$

$$\partial_i c' = \begin{cases} \partial_i c \times N \cup \partial_{j-1} \partial_i c \times I_N & \text{if } i < j, \\ \partial_j c \times 0 & \text{if } i = j, \\ \partial_i c \times N \cup \partial_j \partial_i c \times I_N & \text{if } i > j. \end{cases}$$

So this construction does not change the j -th face $\partial_j c = \partial_j c \times 0$. If $i \neq j$, then one can perform the same construction to $\partial_i c$ to get $(\partial_i c)'$, which coincides with $\partial_i c'$.

Define maps $S_N^v : \Delta \times I_N \rightarrow \Delta \times I_N$ and $\tilde{S}_N^v : M \times I_N \rightarrow M \times I_N$ by

$$S_N^v(x, t) = (s_{t/N}^v(x), t) \quad \text{and} \quad \tilde{S}_N^v(w, t) = (\tilde{s}_{t/N}^v(w), t) .$$

Define an ordered $(n+1)$ -simplex $\Delta^{n+1} (\subset \Delta \times I_N)$ by

$$\Delta^{n+1} = (\langle v_0, \dots, v_j \rangle \times 0) * (\langle v_j \rangle \times N) * (\langle v_{j+1}, \dots, v_n \rangle \times 0) .$$

Here $*$ denotes the join of simplices. Note that

$$S_N^v(\Delta \times I_N) = \bigcup_{0 \leq t \leq N} (s_{t/N}^v(\langle v_0, \dots, v_j \rangle \times t) * (\langle v_{j+1}, \dots, v_n \rangle \times t)) ,$$

$$\Delta^{n+1} = \bigcup_{0 \leq t \leq N} (s_{t/N}^v(\langle v_0, \dots, v_j \rangle \times t) * (\langle v_{j+1}, \dots, v_n \rangle \times 0)) .$$

Therefore, the obvious vertical retraction

$$\langle v_{j+1}, \dots, v_n \rangle \times I_N \longrightarrow \langle v_{j+1}, \dots, v_n \rangle \times 0$$


induces a map R_N^v from the image $S_N^v(\Delta \times I_N)$ to Δ^{n+1} . Let

$$q = p \times 1_{I_N} | : M_{\Delta^{n+1}} = (p \times 1_{I_N})^{-1}(\Delta^{n+1}) \rightarrow \Delta^{n+1}$$

denote the pull-back of $p : M \rightarrow \Delta$ by the projection map

$$\pi : \Delta^{n+1} \xrightarrow{\text{inclusion}} \Delta \times I_N \xrightarrow{\text{projection}} \Delta .$$

The map R_N^v is covered by a map $\tilde{R}_N^v : \tilde{S}_N^v(M \times I_N) \rightarrow M_{\Delta^{n+1}}$.



Let us look at the relation between c and c' (and its functorial image $(\tilde{R}_N^v \circ \tilde{S}_N^v)_*(c')$) more closely. As in the pictures above, define a subset Δ' of $\partial(\Delta \times I_N)$ by

$$\Delta' = \Delta \times N \cup \partial_j \Delta \times I_N .$$

The $(n+2)$ -ad c' lies over Δ' . By glueing some of the faces, let us regard $c \times I_N$ as an $(n+3)$ -ad whose faces are

$$\partial_0 c \times I_N, \dots, \partial_{j-1} c \times I_N, c', c \times 0, \partial_{j+1} c \times I_N, \dots, \partial_n c \times I_N .$$

The functorial image of this $(n+3)$ -ad by the composition $\tilde{R}_N^v \circ \tilde{S}_N^v$ defines a proper quadratic Poincaré $(n+3)$ -ad $\mathcal{C}_N^v(c)$ on $q : M_{\Delta^{n+1}} \rightarrow \Delta^{n+1}$.

The face $(\tilde{R}_N^v \circ \tilde{S}_N^v)_*(c')$ is a proper quadratic Poincaré $(n+2)$ -ad on $q|q^{-1}(R_N^v(S_N^v(\Delta')))$, and is denoted $A_N^v(c)$. Its functorial image $\pi_*(A_N^v(c))$ will be denoted $a_N^v(c)$. It is a proper on Δ . The functorial image $\pi_*(\mathcal{C}_N^v(c))$ can be regarded as a Poincaré cobordism between c and $a_N^v(c)$. The operation described above is called the *Alexander trick (of height N) at the vertex $v = v_j$* . Note that $a_N^v(c)$ has a fine control in the t_j pseudo-coordinate. Also note that $\partial_j a_N^v(c) = a_N^v(\partial_j c) = \partial_j c$, where $v = v_j$.

If we successively apply the Alexander tricks at v_n, \dots, v_1, v_0 to the given proper quadratic Poincaré $(n+2)$ -ad c , then we get finely controlled object which is cobordant to c . This process is called “squeezing” of “shrinking”. When we use the same height N at every vertex, then the squeezed object obtained from c will be denoted $S_N(c)$:

$$S_N(c) = a_N^{v_0}(a_N^{v_1}(\dots(a_N^{v_n}(c))\dots)) .$$

The cobordism between c and $S_N(c)$ constructed above is called the *standard cobordism*. The squeezing operation S_N preserves the face relation:


Proposition 3.1. $\partial_i S_N(c)$ is equal to $S_N(\partial_i c)$. Furthermore, the standard cobordism between $\partial_i c$ and $\partial_i S_N(c)$ is equal to the standard cobordism between $\partial_i c$ and $S_N(\partial_i c)$.

4. \mathbb{L} -SPACES

The squeezing operation seems to justify the following simple definition of the **co-efficient L-space** $\mathbb{L}_n(p : M \rightarrow X)$ for the generalized homology $H_*(X; \mathbb{L}(p))$, where $p : M \rightarrow X$ is a map from a space to a finite polyhedron which has an iterated mapping cylinder decomposition and n is an integer. It is a Δ -set; a k -simplex is an $(n+k)$ -dimensional proper quadratic Poincaré $(k+2)$ -ad $(c; \partial_0 c, \dots, \partial_k c)$ on the pull-back $\pi^* M \rightarrow (\Delta; \partial_0 \Delta, \dots, \partial_k \Delta)$, where Δ is a k -simplex and $\pi : \Delta \rightarrow \Delta^l$ is an affine surjection from Δ to an l -dimensional simplex Δ^l of X ($l \leq k$) induced by an order(\leq) preserving map between the vertices.


Two such simplices $(c, \pi : \Delta \rightarrow \Delta^l)$ and $(c', \pi' : \Delta' \rightarrow \Delta^l)$ are identified when there is an affine homeomorphism $\phi : \Delta \rightarrow \Delta'$ of ordered simplices such that $\pi = \pi' \circ \phi$ and $\phi_*(c) = c'$.

Note that the squeezing operation S_N defines a simplicial homotopy of the identity map of $\mathbb{L}_n(p : M \rightarrow X)$ to a simplicial map whose image is contained in a subset made up of simplices of ‘small radius’ measured on X , if N is large. Thus this space has a built-in ‘squeezing’ mechanism.



Let us consider the special case when X is a single point. There is a similar Δ -set $\mathbb{L}'_n(M)$ whose k -simplex is an $(n+k)$ -dimensional quadratic Poincaré $(k+2)$ -ad c on M that is *special*, i.e. $\partial_0 \partial_1 \dots \partial_k c$ is 0. $\pi_0(\mathbb{L}'_n(p : M \rightarrow *))$ is isomorphic to $L_n^h(\mathbb{Z}\pi_1(M))$.

There is a map $\mathbb{L}_n(M \rightarrow *) \rightarrow \mathbb{L}'_n(M)$ that sends a k -simplex (c, π) to its functorial image $\pi_*(c)$. A map in the reverse direction can be constructed as follows. Let c be a k -simplex of $\mathbb{L}'_n(M)$. It is made up of three type of things: (1) ‘points’ in M (generators of free modules), (2) paths with coefficients connecting the generators, and (3) homotopies of certain paths. Since c is special, one can make a 1–1 correspondence between its faces (including c itself) and the faces of a standard k -simplex Δ (including Δ itself), and can make copies of the faces of c on the sets $\{\text{barycenters}\} \times M \subset \Delta \times M$ and realizing the morphisms between adjacent pieces by using the original paths in c in the M -direction and the path connecting two adjacent barycenters in the Δ -direction as components. Similarly for homotopies of paths. These are homotopy inverses of each other.



Therefore, $\mathbb{L}_n(p : M \rightarrow X)$ defined above may give a convenient description of \mathbb{L} -homology groups.

REFERENCES

1. A. E. Hatcher, *Higher simple homotopy theory*, Ann. of Math., **102** (1975) 101 – 137
2. E. K. Pedersen and M. Yamasaki, *Stability in Controlled L-theory*, in *Exotic homology manifolds – Oberwolfach 2003 (electronic)*, Geom. Topol. Monogr. **9** (Geom. Topol. Publ., Coventry, 2006) 67 – 86.
3. C. P. Rourke and B. J. Sanderson, *Δ -sets I: homotopy theory*, Quart. J. Math. **22** (1971) 321 – 338.
4. M. Yamasaki, *L-groups of crystallographic groups*, Invent. Math. **88** (1987) 571 – 602
5. M. Yamasaki, *L-groups of virtually polycyclic groups*, Topology Appl. **33** (1989) 223 – 233